

FIELD 2007



Southwest Research-Extension Center



**Report of Progress
980**

*Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service*

KANSAS STATE

Southwest Research-Extension Center

EFFICACY OF FIPRONIL APPLIED AS A FOLIAR TREATMENT ON SIX COMMERCIAL SOYBEAN VARIETIES TO CONTROL DECTES STEM BORERS IN SOYBEAN, SCANDIA, KS, 2006

by

Larry Buschman, Teru taka Niide¹, William Schapaugh², and Barney Gorden³

SUMMARY

We tested a foliar fipronil insecticide treatment applied to six soybean varieties to determine effectiveness in reducing *Dectes* stem borers (*Dectes texanus*) in soybean. The foliar application of fipronil significantly reduced *Dectes* stem borer infestations between 76% and 88%. However, these treatments increased yield only 2.9%, and this was not statistically significant. *Dectes* stem borer infestation averaged 55% infested plants.

PROCEDURES

Seed of six commercial soybean varieties in maturity groups II through to IV was machine-planted at 16 seed per row-foot on May 17, 2005, at the irrigation experiment field near Scandia,. The plots were four rows wide and 20 feet long. There was a 3-foot-wide alley at each end of the plot. The design was a randomized block experiment with three replications. There was a treated and untreated plot of each variety in each replication. The foliar treatment of fipronil was applied July 18 during the beetle flight. This treatment targeted the first two instars developing inside the plants. The foliar treatment was applied with a backpack sprayer, using a hand-held boom with two nozzles (Conejet TXVS 6) directed at a single row. The nozzles were held 6-8 inches from the plants to maximize coverage of the upper canopy. The sprayer was calibrated to deliver 20 gal/acre (8.0 sec per 20 ft row at 30 psi). A chronometer was used to measure the time spent on each row to help maintain appropriate speed.

The experiment was analyzed as a two-factor experiment with six levels of variety and two levels of treatment.

Dectes stem borer infestations were recorded at the end of the season (September 22) by dissecting five consecutive plants from each of the four rows in each plot for a total of 20 plants. The plants were dissected to record entry nodes, upper stem tunneling, tunneling that reached the base of the plant, and presence of live *Dectes* larvae. Grain yield data was collected by machine harvesting the plots October 12 and converted to bushels per acre based on 12% moisture.

RESULTS AND DISCUSSION


Dectes stem borer infested 55% of plants in 2006. This was similar to the infestation of 2004, when we were able to show yield responses with fipronil treatment. In this 2006 trial, the fipronil treatment significantly reduced *Dectes* stem borer infestations (81%, 78%, 82% and 88% for entry nodes, stem tunneling, base tunneling and live larvae, respectively; Table 1). However, the fipronil treatment only increased grain yield 2.9%, and this was not a significant increase. There were some significant differences in *Dectes* infestation across the different varieties, but there was no significant yield difference across the varieties. This was surprising because there was such a wide difference in maturity across the varieties. The 2006 results suggest there was no physiological yield loss associated with *Dectes* stem borer infestations. We were not able to show differences in tolerance of the different varieties to *Dectes* stem borer infestations.

¹Department of Entomology, Kansas State University, Manhattan

²Department of Agronomy, Kansas State University, Manhattan

³Department of Irrigation and North Central Kansas Experimental Fields, Kansas State University, Scandia

Dectes Stem Borer Entry Nodes per 20 Plants




Variety P=0.0027
Treatment P=>0.0001

Soybean Variety

■ UnSprayed ■ Sprayed

81% Control

Dectes Stem Borer Larvae per 20 Plants




Variety P=<0.5000
Treatment P=>0.0001

Soybean Variety

■ UnSprayed ■ Sprayed

88% Control

Grain Yield Bu/Acre Sprayed and Unsprayed Soybean Varieties



Variety P= 0.3909
Treatment P=<0.5000

Soybean Variety

■ UnSprayed ■ Sprayed

2.9% Increase

Table 1. F-test Probability values for the ANOVA tests of the two main effects, variety and insecticide treatment. Fipronil treatments were applied as foliar treatments. Irrigation Experiment Field, Scandia, Kansas, 2006.

	Soybean Maturity Group	Entry Nodes /20 plants	Stem Tunneling /20 plants	Base Tunneling /20 plants	Live Larvae /20 plants	Grain Yield Bu/Acre
ANOVA F-Test Probability						
Replication		0.001	0.0191	0.0669	0.0566	0.0383
Variety		0.0027	0.3505	0.0087	<0.5000	0.3909
Insecticide		>0.0001	>0.0001	>0.0001	>0.0001	<0.5000
V x I Interaction		0.0175	<0.5000	0.0465	<0.5000	<0.5000
Variety Means—Untreated						
Nex2403K2RR	Mid II	8.7	7.3	4.7	4.3	58.9
Dyna-GroDB32C25	Early III	25.0	12.7	2.3	7.7	61.4
Pioneer 93M50	Mid III	17.7	13.3	2.0	6.0	69.0
Ohlde 3727NRS	Late III	13.7	10.3	8.0	6.7	63.2
KS4404RR	Early IV	16.7	12.0	6.0	3.3	69.7
KS4704RR	Mid IV	13.0	9.7	5.3	5.7	66.9
Mean		15.8	10.9	4.7	5.6	64.85
Variety Means—Fipronil—Treated						
Nex2403K2RR	Mid II	2.7	2.7	1.0	0.7	67.5
Dyna-GroDB32C25	Early III	4.0	3.0	1.0	0.7	61.4
Pioneer 93M50	Mid III	4.0	3.0	0.0	0.0	69.0
Ohlde 3727NRS	Late III	1.7	1.0	1.0	0.7	63.2
KS4404RR	Early IV	2.7	2.7	1.0	1.0	69.7
KS4704RR	Mid IV	3.3	2.3	1.0	1.0	66.9
Mean		3.06	2.5	0.8	0.7	66.76

Note: Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned.

This report was produced by the Department of Communications at Kansas State University. These materials may be freely reproduced for educational purposes. All other rights reserved. In each case, give credit to the author(s), name of work, Kansas State University, and the date the work was published.

Kansas State University Agricultural Experiment Station and Cooperative Extension Service, Manhattan 66506

SRP 980

August 2007

K-State Research and Extension is an equal opportunity provider and employer. These materials may be available in alternative formats.

300